



## Prevalence of musculoskeletal symptoms in blue-collar workers: association with individual and lifestyle related factors

Isabel Moreira-Silva<sup>a</sup>, Joana Azevedo<sup>b</sup>, Sandra Rodrigues<sup>c</sup>, Nuno Ventura<sup>d</sup>, Adérito Seixas<sup>e</sup>,  
Jorge Mota<sup>f</sup>

<sup>a</sup>Faculty of Health Sciences, University Fernando Pessoa, Porto, PT and LABIOMPEP, INEGI-LAETA, Faculty of Sports, University of Porto, PT ([isabelmsilva@ufp.edu.pt](mailto:isabelmsilva@ufp.edu.pt)) ORCID: 0000-0002-4137-7694. <sup>b</sup>Faculty of Health Sciences, University Fernando Pessoa, Porto, PT ([jsazevedo@ufp.edu.pt](mailto:jsazevedo@ufp.edu.pt)) ORCID: 0000-0002-3616-8679. <sup>c</sup>Faculty of Health Sciences, University Fernando Pessoa, Porto, PT ([sandrar@ufp.edu.pt](mailto:sandrar@ufp.edu.pt)) ORCID: 0000-0003-2931-8971. <sup>d</sup>Faculty of Health Sciences, University Fernando Pessoa, Porto, PT ([nunov@ufp.edu.pt](mailto:nunov@ufp.edu.pt)) ORCID: 0000-0003-2104-2480. <sup>e</sup>Faculty of Health Sciences, University Fernando Pessoa, Porto, PT and LABIOMPEP, INEGI-LAETA, Faculty of Sports, University of Porto, PT ([aderito@ufp.edu.pt](mailto:aderito@ufp.edu.pt)) ORCID: 0000-0002-6563-8246. <sup>f</sup>CIAFEL, Faculty of Sports, University of Porto, Porto, PT ([jmota@fade.up.pt](mailto:jmota@fade.up.pt)) ORCID: 0000-0001-7571-9181.

### Article History

Received 2 July 2019  
Accepted 24 October 2019  
Published 15 November 2019

### Keywords

Musculoskeletal symptoms  
Blue-collar workers  
Individual factors  
Lifestyle factors

### DOI:

10.24840/2184-0954\_003.003\_0001

### ISSN:

2184-0954

### Type:

Research Article



Open Access



Peer Reviewed



CC BY

### Abstract

This study aimed to investigate the 12-month and 7-day prevalence of musculoskeletal symptoms in different body regions and its association with different individual and lifestyle-related factors among blue-collar workers of a Portuguese manufacturing company. One hundred and thirty-six blue-collar workers participated in the study. Musculoskeletal symptoms were assessed using the Nordic Musculoskeletal Questionnaire, and the physical activity level and sitting time were assessed with the IPAQ-Short Version. The 12-month prevalence was higher in the low back region (56.6%), followed by the wrist/hand (50%), the shoulder (45.6%) and the neck (44.9%). In the last 7 days, the four most affected body regions were: the low back region (25%), the shoulder (20.6%) and the neck and wrist/hand (19.9%). Regarding individual factors, significant associations were found between age and the prevalence of musculoskeletal symptoms in the shoulder ( $\chi^2(2)=6.783$ ;  $p=0.034$ ), elbow ( $\chi^2(2)=6.813$ ;  $p=0.033$ ), wrist/hand ( $\chi^2(2)=6.999$ ;  $p=0.030$ ), thigh/hip ( $\chi^2(2)=9.771$ ;  $p=0.008$ ), neck ( $\chi^2(2)=9.293$ ;  $p=0.010$ ) and the low back region ( $\chi^2(2)=6.190$ ;  $p=0.045$ ), with the older workers reporting a higher prevalence of musculoskeletal symptoms. Moreover, women reported a significantly higher prevalence of musculoskeletal symptoms in the neck than men ( $\chi^2(1)=5.054$ ;  $p=0.025$ ). Overweight and obese workers had a significantly higher prevalence of symptoms in the shoulder ( $\chi^2(2)=10.595$ ;  $p=0.005$ ) and neck ( $\chi^2(2)=6.521$ ;  $p=0.038$ ). Regarding lifestyle-related factors, no significant associations were found between physical activity level or sitting time ( $p>0.05$ ) and the prevalence of musculoskeletal symptoms in any of the body regions. To conclude, blue-collar workers have a high prevalence of musculoskeletal symptoms. Individual factors like age, gender and BMI seem to influence the prevalence of musculoskeletal symptoms in different body regions, but lifestyle-related factors such as the physical activity level and sitting time seem not to be associated with the prevalence of musculoskeletal symptoms in the blue-collar workers of this sample. The present results emphasize the need of workplace interventions to prevent musculoskeletal symptoms in this population.

## 1. INTRODUCTION

The sectorization of labor, that emerged from the Industrial Revolution, forced workers to perform tasks repetitively or to stay for longer periods of time in a maintained posture during their working hours, which causes pain, physical discomfort and, consequently, musculoskeletal disorders (Gupta *et al.*, 2016; Sjögrena *et al.*, 2006). Musculoskeletal disorders describe a group of injuries and/or disabilities affecting muscles, bones, nerves, tendons, ligaments, joints, cartilages or spinal discs (Bureau of Labor Statistics Occupational Safety and Health, 2012). According to the World Health Organization (WHO), work-related musculoskeletal disorders have multifactorial causes or risk factors (WHO, 1995).

Blue-collar workers frequently represent an occupational category with higher physical workloads, spending most of their labor time standing or walking (Chen, Perry, Yang, & Yang, 2017; Schreuder, Roelen, Koopmans, & Groothoff, 2008; Yildirim, Gunay, & Karadibak, 2014). According to different authors, there is an annual prevalence of musculoskeletal symptoms in these employees ranging from 61% (Aghilinejad, Choobineh, Sadeghi, Nouri, & Ahmadi, 2012) to 72% (Ghasemkhani, Mahmudi, & Jabbari, 2008). Likewise, it is stated that workers engaging higher physical workloads report a higher prevalence of symptoms and in more body regions than those with lighter workloads (Widanarko *et al.*, 2011). Furthermore, the high prevalence of musculoskeletal disorders in blue-collar workers is a major cause of absenteeism, diminished working abilities, higher medical costs and decreased productivity (Hanebuth, Meinel, & Fischer, 2006; Manjunatha, Kiran, & Thankappan, 2011).

The contribution of the individual factors to the prevalence of musculoskeletal symptoms, such as age, gender or anthropometric characteristics (e.g. Body Mass Index) have been widely investigated. Regarding age, most of the available literature reported that older workers in physically demanding jobs seem to complaint more about musculoskeletal disorders than younger employees (de Zwart, Broersen, Frings-Dresen, & Van Dijk, 1997; Hussain, 2004). Similarly, when compared to men, women often report a higher number of painful regions and a higher prevalence of musculoskeletal symptoms (de Zwart *et al.*, 1997; Widanarko *et al.*, 2011). Other authors also reported a significant association between Body Mass Index (BMI) and the presence of musculoskeletal symptoms, especially in the lower limbs (Andersen, Haahr, & Frost, 2007), with overweight and obese workers reporting higher 12-month prevalence of musculoskeletal symptoms and having a higher probability of developing work-related musculoskeletal disorders (Aghilinejad *et al.*, 2012; Viester *et al.*, 2013).

Regarding the influence of lifestyle-related factors, it has been stated that blue-collar workers submitted to prolonged sitting during their labor hours are more likely to develop musculoskeletal disorders (Hallman, Gupta, Mathiassen, & Holtermann, 2015). Furthermore, Van Amelsvoort, Spigt, Swaen, and Kant (2006) demonstrated a significantly decrease in sick leaves caused by musculoskeletal disorders when workers were active in their leisure time, compared to workers with a more sedentary behavior, which suggests that physical activity is beneficial. Similarly, several authors demonstrated positive effects of different workplace physical activity interventions on sick leave, costs, new episodes of pain, physical discomfort, and consequently, musculoskeletal disorders among workers (Kellett, Kellett, & Nordholm, 1991; Kietrys, Galper, & Verno, 2007; Tveito, Hysing, & Eriksen, 2004; Tveito & Eriksen, 2009; Zebis *et al.*, 2011).

Furthermore, other risk factors should be recognized. Especially in the low back region, different investigations recognized occupational risk factors such as inadequate standing and sitting postures, heavy lifting and repetitive movements, to be significant risk factors for pain in this body region among different industrial workers (Goswami *et al.*, 2016; Miranda, Viikari-Juntura, Punnett, & Riihimaki, 2008; Yildirim *et al.*, 2014). Similarly, risk factors associated with the psychosocial context of work, namely poor workplace social environment, job dissatisfaction or low job control, have been shown to influence pain and other symptoms in the low back region of workers of an automobile manufacturing company (Kerr *et al.*, 2001). Likewise, a study conducted with offshore workers from an oil company demonstrated that occupational stressors, in particular stress with their own safety, were important predictors of musculoskeletal pain in different body regions, highlighting the role of psychosocial factors in work-related musculoskeletal disorders, often little taken into consideration (Chen, Yu, & Wong, 2005).

Workplace health promotion initiatives are programs that aim to promote health by reducing its risks and actively prevent the beginning of disease (Goetzel & Ozminkowski, 2008). This may be operationalized using different types of strategies, namely decreasing stress, excess of body weight, improving nutrition and increasing physical activity and fitness levels. Indeed, in the last 20 years, the number of these programs has been growing, as their advantages for employees and companies are increasingly being recognized, as they help to reduce employee's health care costs and disability and improve employee's productivity (Wang *et al.*, 2010).

Studies in different countries have investigated the prevalence of musculoskeletal symptoms in manufacturing or industrial workers like blue-collar workers and the association with different risk factors. However, according to Serranheira (2007), in Portugal, the limited data available reported by the National Centre for Protection Against Occupational Hazards revealed a gradual increase in the number of cases of musculoskeletal disorders and the consequent increase of absenteeism and the use of the Portuguese health care system and health insurance companies by workers. According to the same author there is a lack of studies investigating this matter,

especially the 7-day prevalence, and its association with different factors in this particular occupational group.

In that sense, the aim of the present study was to investigate the 12-month and 7-day prevalence of musculoskeletal symptoms in different body regions and its association with different individual (age, gender and BMI) and lifestyle-related factors (physical activity level and sitting time) among blue-collar workers of a Portuguese manufacturing company.

## **2. MATERIALS AND METHODS**

### **2.1. Study Design and sampling**

This observational, cross-sectional study was approved by the ethical committee of the local University and by the board of the company. The study was conducted in a Portuguese multinational manufacturing company of water heaters. There are 1000 employees in the company, however, only a convenience sample of 220 were allowed by the administration board to participate. These employees perform prolonged shifts (7 hours and 30 minutes) and repetitive demanding tasks using mainly the upper limbs, while in the standing position. The main tasks that these workers perform during their working hours are loading heavy objects, use screwdrivers and place washers in equipment such as water heaters.

All the participants were full-time workers (40h/week) and had been employed in the company for at least 6 months. From the 220 employees invited, 209 agreed to participate (87 male; 122 female), with 136 of them being considered blue-collar workers (40 male; 96 female). The mean age of these workers was  $37.4 \pm 8.5$  years and the mean BMI was  $27.2 \pm 4.9$  Kg/m<sup>2</sup>, which according to the World Health Organization (2019) represents a pre-obesity category. All participants expressed their consent to participate in the study by signing an informed consent form.

### **2.2. Individual data collection**

Individual factors such as age and gender were self-reported. Height (m) of the participants was measured with the participants standing upright against a stadiometer in bare or stocking feet (Holtain Ltd., Crymmych, Pembrokeshire, UK). Weight (kg) was measured using a portable electronic weight scale (Tanita Inner Scan BC 532, Tokyo, Japan). BMI was calculated from the ratio between body weight and body height (kg/m<sup>2</sup>). Participants BMI was expressed as normal (18.5-24.9), overweight (25.0-29.9) and obese (>30.0), according to the World Health Organization (2019).

### **2.3. Musculoskeletal disorders and related symptoms**

Musculoskeletal pain and related symptoms were assessed with the Standardized Nordic Musculoskeletal Questionnaire (NMQ) (Kuorinka, 1987), supplemented with questions about localized pain intensity. This questionnaire has already been validated to the Portuguese population by Mesquita, Ribeiro, and Moreira (2010). The NMQ consists of 27 questions of binary choice (yes or no).

The questionnaire has three questions correlating to nine anatomic regions (neck, shoulders, elbows, wrists/hands, dorsal region, low back region, hips/thighs, knees and ankles/feet). The first is "had some troubles or pain in the last 12 months"; the second is "in the last 12 months did you feel some limitation caused by work in the daily activities"; and the third is "had some troubles or pain in the last 7 days" (Kuorinka, 1987). The questionnaire also includes the numeric pain scale (scale 0-10) to assess the pain intensity in the last 7 days.

### **2.4. Physical Activity Level and Sitting Time Assessment**

Physical activity level and sitting time were assessed using the short version of the International Physical Activity Questionnaire (IPAQ) (Craig *et al.*, 2003). According to the Guidelines for the data processing and analysis of the IPAQ, total physical activity was expressed as metabolic equivalent (MET) minutes/week by weighting the reported minutes per week in each activity category by the metabolic equivalent specific to each activity (Total Physical Activity = 3.3 MET x walking minutes x walking days + 4.0 MET x moderate-intensity activity minutes x moderate days + 8.0 MET x vigorous-intensity activity minutes x vigorous-intensity days). Physical activity level of the participants was expressed as active, inactive and insufficiently active. Regarding the sitting time, blue-collar workers of this sample spend a median of 120 minutes sitting during

the day.

## 2.5. Statistical Analysis

Data was analyzed with the Statistical Package for the Social Sciences (SPSS v. 25.0) software for Windows. Descriptive characteristics of the participants (age and BMI) were presented as mean  $\pm$  standard deviation. The prevalence of musculoskeletal symptoms during the last 12 months and the last 7 days in each body region was described in percentage (%). The Chi-Square test was used to assess the association between the prevalence of musculoskeletal symptoms and the individual (age, gender and BMI) and lifestyle-related factors (physical activity level and sitting time). A p value under 0.05 was considered statistically significant.

## 3. RESULTS

### 3.1. 12-month and 7-day prevalence of musculoskeletal symptoms in blue-collar workers

The analysis of the NMQ data demonstrated that the 12-month prevalence of musculoskeletal symptoms is high, especially in the low back region (56.6%), followed by the wrist/hand (50%), the shoulder (45.6%) and the neck (44.9%). In the other body regions, the prevalence was 30.1% in the ankle/feet; 22.8% in the knee; and 19.1% in the elbow and thigh/hip regions.

Regarding the 7-day prevalence, the most affected body regions were: the low back region (25%), followed by the shoulder (20.6%), the wrist/hand and the neck (19.9%). In the other body regions, the prevalence was 13.2% in the ankles/feet; 8.1% in the elbow and knee; and 7.4% in the thigh/hip.

In tables 1 to 5, the percentage represents the proportion of participants reporting pain in each body region by category (column). It is also important to note that each participant may report pain in more than one body region.

### 3.2. Association between musculoskeletal symptoms and individual factors (age, gender and BMI)

Significant associations were found between age and the prevalence of musculoskeletal symptoms in the shoulder ( $p=0.034$ ), elbow ( $p=0.033$ ), wrist/hand ( $p=0.030$ ), thigh/hip ( $p=0.008$ ), neck ( $p=0.010$ ) and the low back region ( $p=0.045$ ), with the older workers within the 41-60 age group presenting a higher prevalence of musculoskeletal symptoms, except for the low back region in which the highest prevalence was found in the 31-40 age group (Table 1).

**Table 1.** Association between musculoskeletal symptoms and age

Body Regions	20-30 years (n=33)	31-40 years (n=59)	41-60 years (n=44)	Chi-Square Test
Shoulder (n=62)	9 (27.3%)	28 (47.5%)	25 (56.8%)	$\chi^2(2)=6.783$ ; $p=0.034^*$
Elbow (n=26)	4 (12.1%)	8 (13.6%)	14 (31.8%)	$\chi^2(2)=6.813$ ; $p=0.033^*$
Wrist/Hand (n=68)	10 (30.3%)	32 (54.2%)	26 (59.1%)	$\chi^2(2)=6.999$ ; $p=0.030^*$
Thigh/Hip (n=26)	5 (15.2%)	6 (10.2%)	15 (34.1%)	$\chi^2(2)=9.771$ ; $p=0.008^*$
Knee (n=31)	5 (15.2%)	12 (20.3%)	14 (31.8%)	$\chi^2(2)=3.333$ ; $p=0.189$
Ankles/Feet (n=41)	7 (21.2%)	22 (37.3%)	12 (27.3%)	$\chi^2(2)=2.852$ ; $p=0.240$
Neck (n=61)	8 (24.2%)	27 (45.8%)	26 (59.1%)	$\chi^2(2)=9.293$ ; $p=0.010^*$
Dorsal region (n=19)	5 (15.2%)	8 (13.6%)	6 (13.6%)	$\chi^2(2)=0.051$ ; $p=0.975$
Low back region (n=77)	14 (42.4%)	32 (55.2%)	31 (23.0%)	$\chi^2(2)=6.190$ ; $p=0.045^*$

\* $p<0.05$

Gender was significantly associated with the prevalence of musculoskeletal symptoms in the neck region ( $p=0.025$ ), with females presenting a higher prevalence than male workers (Table 2).

**Table 2.** Association between musculoskeletal symptoms and gender

Body Regions	Male (n=40)	Female (n=96)	Chi-Square Test
Shoulder (n=62)	16 (40.0%)	46 (47.9%)	$\chi^2(1) = 0.713$ ; $p=0.398$
Elbow (n=26)	4 (10.0%)	22 (22.9%)	$\chi^2(1) = 3.047$ ; $p=0.081$
Wrist/Hand (n=68)	16 (40.0%)	52 (54.2%)	$\chi^2(1) = 2.267$ ; $p=0.132$
Thigh/hip (n=26)	6 (15.0%)	20 (20.8%)	$\chi^2(1) = 0.621$ ; $p=0.431$
Knee (n=31)	10 (25.0%)	21 (21.9%)	$\chi^2(1) = 0.157$ ; $p=0.692$
Ankles/Feet (n=41)	13 (32.5%)	28 (29.2%)	$\chi^2(1) = 0.149$ ; $p=0.700$
Neck (n=61)	12 (30.0%)	49 (51.0%)	$\chi^2(1) = 5.054$ ; <b><math>p=0.025^*</math></b>
Dorsal Region (n=19)	5 (12.5%)	14 (14.6%)	$\chi^2(1) = 0.102$ ; $p=0.749$
Low back region (n=77)	20 (51.3%)	57 (59.4%)	$\chi^2(1) = 0.741$ ; $p=0.389$

\* $p<0.05$ 

Significant associations were found between BMI and the prevalence of musculoskeletal symptoms. Obese workers had a significantly higher prevalence of musculoskeletal symptoms in the neck ( $p=0.038$ ) in comparison with the other BMI categories, and overweight workers had a significantly higher prevalence of symptoms in the shoulder ( $p=0.005$ ) (Table 3).

**Table 3.** Association between musculoskeletal symptoms and BMI (in kg/cm<sup>2</sup>)

Body Regions	Normal (n=53)	Overweight (n=48)	Obese (n=35)	Chi-Square Test
Shoulder (n=62)	15 (28.3%)	28 (58.3%)	19 (54.3%)	$\chi^2(2)=10.595$ ; <b><math>p=0.005^*</math></b>
Elbow (n=26)	7 (13.2%)	11 (22.9%)	8 (22.9%)	$\chi^2(2)=1.962$ ; $p=0.375$
Wrist/Hand (n=68)	22 (41.5%)	24 (50.0%)	22 (62.9%)	$\chi^2(2)=3.843$ ; $p=0.146$
Thigh/Hip (n=26)	6 (11.3%)	12 (25.0%)	8 (22.9%)	$\chi^2(2)=3.474$ ; $p=0.176$
Knee (n=31)	10 (18.9%)	8 (16.7%)	13 (37.1%)	$\chi^2(2)=5.583$ ; $p=0.061$
Ankles/Feet (n=41)	14 (26.4%)	13 (27.1%)	14 (40.0%)	$\chi^2(2)=2.178$ ; $p=0.337$
Neck (n=61)	19 (35.8%)	20 (41.7%)	22 (62.9%)	$\chi^2(2)=6.521$ ; <b><math>p=0.038^*</math></b>
Dorsal region (n=19)	11 (20.8%)	4 (8.3%)	4 (11.4%)	$\chi^2(2)=3.487$ ; $p=0.175$
Low back region (n=77)	29 (54.7%)	26 (57.4%)	21 (60.0%)	$\chi^2(2)=0.245$ ; $p=0.885$

\* $p<0.05$ 

### 3.3. Association between musculoskeletal symptoms and lifestyle-related factors (physical activity level and sitting time)

No significant associations were found between physical activity level (Table 4) or sitting time (Table 5) and the prevalence of musculoskeletal symptoms in the different body regions ( $p>0.05$ ).

**Table 4.** Association between musculoskeletal symptoms and physical activity level

Body Regions	Active (n=104)	Inactive (n=19)	Insufficiently Active (n=13)	Chi-Square Test
Shoulder (n=62)	51 (49.0%)	5 (26.3%)	6 (46.2%)	$\chi^2(2) = 3.346$ ; $p=0.188$
Elbow (n=26)	19 (18.3%)	4 (21.1%)	3(23.1%)	$\chi^2(2) = 0.226$ ; $p=0.893$
Wrist/Hand (n=68)	53 (51.0%)	7 (36.8%)	8(61.5%)	$\chi^2(2) = 2.047$ ; $p=0.359$
Thigh/hip (n=26)	22 (21.2%)	4 (21.1%)	0 (0%)	$\chi^2(2) = 3.398$ ; $p=0.183$
Knee (n=31)	26 (25.0%)	4 (21.1%)	1(7.7%)	$\chi^2(2) = 2.005$ ; $p=0.367$
Ankles/Feet (n=41)	30 (28.8%)	5 (26.3%)	6 (46.2%)	$\chi^2(2) = 1.798$ ; $p=0.407$
Neck (n=61)	45 (43.3%)	9 (47.4%)	7 (53.8%)	$\chi^2(2) = 0.579$ ; $p=0.749$
Dorsal Region (n=19)	17 (16.3%)	1 (5.3%)	1 (7.7%)	$\chi^2(2) = 2.113$ ; $p=0.348$
Low back region (n=77)	61 (59.2%)	7 (36.8%)	9 (69.2%)	$\chi^2(2) = 4.152$ ; $p=0.125$

**Table 5.** Association between musculoskeletal symptoms and sitting time (in minutes)

Body Regions	<120 min (n=58)	≥120 min (n=78)	Chi-Square Test
Shoulder (n=62)	28 (48.3%)	34 (43.6%)	$\chi^2(1) = 0.294$ ; $p=0.587$
Elbow (n=26)	14 (24.1%)	12 (15.4%)	$\chi^2(1) = 1.648$ ; $p=0.199$
Wrist/Hand (n=68)	33 (56.9%)	35 (44.9%)	$\chi^2(1) = 1.924$ ; $p=0.165$
Thigh/hip (n=26)	9 (15.5%)	17 (21.8%)	$\chi^2(1) = 0.848$ ; $p=0.357$
Knee (n=31)	11 (19.0%)	20 (25.6%)	$\chi^2(1) = 0.842$ ; $p=0.359$
Ankles/Feet (n=41)	15 (25.9%)	26 (33.3%)	$\chi^2(1) = 0.882$ ; $p=0.348$
Neck (n=61)	26 (44.8%)	35 (44.9%)	$\chi^2(1) = 0.000$ ; $p=0.996$
Dorsal Region (n=19)	5 (8.6%)	14 (17.9%)	$\chi^2(1) = 2.408$ ; $p=0.121$
Low back region (n=77)	33 (56.9%)	44 (57.1%)	$\chi^2(1) = 0.001$ ; $p=0.977$

When analyzing the association between reported pain and sitting time, regardless of the body region, no significant association was found (Table 6).

**Table 6.** Association between musculoskeletal symptoms, regardless of body region, and sitting time (in minutes)

Body Regions	<120 min	≥120 min	Chi-Square Test
Pain	49 (36.0%)	67 (49.3%)	$\chi^2(1) = 0.053; p=0.818$
No Pain	9 (6.6%)	11 (8.1%)	

#### 4. DISCUSSION

This study investigated the 12-month and 7-day prevalence of musculoskeletal symptoms in different body regions and its association with different individual (age, gender and BMI) and lifestyle-related factors (physical activity level and sitting time) among blue-collar workers of a Portuguese manufacturing company.

The results of the present study revealed that the prevalence of musculoskeletal symptoms is high in blue-collar workers, with the four most affected body regions in the last 12 months being the low back region (56.6%), followed by the wrist/hand (50%), the shoulder (45.6%) and the neck (44.9%). Regarding the 7-day prevalence, the percentages are lower, even though the four most affected body regions are the same, with the low back region similarly registering the highest prevalence (25%), followed by the shoulder (20.6%) and the wrist/hand and neck with 19.9%.

The majority of investigations in blue-collar workers also confirm that the low back seems to be the body region where the highest prevalence of pain is described (Ng, Tamrin, Yik, Yusoff, & Mori, 2013; Widanarko, Legg, Stevenson, Devereux, & Jones, 2013). Previous studies have reported a prevalence of 54% (Widanarko et al., 2013) and 58% (Ng et al., 2013), which is very close to the percentages reported here, regarding the 12-month prevalence. Indeed, the workers of this sample perform heavy loading and lifting tasks during their working hours, which according to Miranda et al. (2008) are significant predictors of low back pain among blue-collar workers. Regions such as the neck and the shoulder are also pointed out as the most prevalent regions with musculoskeletal symptoms in this population, registering in previous investigations a prevalence of 43% and 42%, respectively, which is also very close with the results presented in this paper (Widanarko et al., 2013).

The high prevalence of musculoskeletal disorders in this group of workers may be related to the high physical demands. Effectively, blue-collar workers, as previously mentioned, represent an occupational group with higher physical workloads (Chen et al., 2017; Schreuder et al., 2008; Yildirim et al., 2014). When physical work demands exceed the safety margin of the individual physical capacities, the risk of physical deterioration increases and can be translated into musculoskeletal disorders, poor work ability and sickness absenteeism (Holtermann et al., 2010). However, workload was not assessed objectively in this study and a causative relationship cannot be established.

As previously described, the contribution of different individual factors to the prevalence of musculoskeletal pain and related symptoms have been widely investigated in the literature. In the present investigation, significant associations were found between age, gender and BMI and the prevalence of musculoskeletal symptoms in different body regions.

Regarding age, the older blue-collar workers of this sample demonstrated a higher prevalence of musculoskeletal symptoms. Indeed, similar results were reported by de Zwart et al. (1997), demonstrating that musculoskeletal complaints among workers in a physically demanding job increases with age, which can be a consequence of the physiological degenerative processes associated with aging (De Magalhães, Curado, & Church, 2009). Furthermore, a higher number of years in the same job continually exposes the worker to the same occupational risk factors, leading to higher or continuous musculoskeletal complaints (Tinubu, Mbada, Oyeyemi, & Fabunmi, 2010).

The results of this study also revealed an association between gender and the prevalence of musculoskeletal symptoms although only in the neck, with women reporting a significantly higher prevalence than male workers of this manufacturing company. Overall, for the other body regions, the present study also found a tendency for a higher prevalence of symptoms in females, except for the knee and ankles/feet. Previous investigations in workers with physically demanding occupations have also reported significant differences between male and female gender, with most of them similarly confirming a higher prevalence of neck symptoms in females,

corroborating our results (Miranda, Viikari-Juntura, Martikainen, Takala, & Riihimäki, 2001; Widanarko *et al.*, 2011). However, these authors also reported significant associations between gender and musculoskeletal symptoms in other body regions, such as the shoulders, the wrists/hands, the upper back and the hips/thighs, with women also reporting higher prevalence of symptoms and in more body regions than male employees (Widanarko *et al.*, 2011).

The present results also revealed that obese workers had a significantly higher prevalence of musculoskeletal symptoms in the neck, and that overweight workers had a significantly higher prevalence of symptoms in the shoulder, when compared to the ones within the normal BMI category. Results of previous investigations have similarly suggested that overweight and obese workers have higher 12-month prevalence of musculoskeletal symptoms, being at a higher probability of developing work-related musculoskeletal disorders (Aghilinejad *et al.*, 2012; Viester *et al.*, 2013). However, in the studies conducted by Andersen *et al.* (2007) and Viester *et al.* (2013), significant associations between BMI and musculoskeletal symptoms were reported essentially in the lower limbs (Andersen *et al.*, 2007; Viester *et al.*, 2013), unlike our results that reported significant associations essentially in the upper limbs. These differences in results may be attributable to the specificities of the studied samples. Both studies (Andersen *et al.*, 2007; Viester *et al.*, 2013) analysed general working populations and the present study analysed workers using mainly the upper limbs, which may have influenced the results.

In order to verify the influence of the lifestyle-related factors on the prevalence of musculoskeletal symptoms, this study conducted both an analysis of the physical activity level of the participants as well as the time they spend in the sitting position.

In blue-collar workers, it has been described that prolonged sitting during the working hours leads to musculoskeletal disorders, particularly in the neck and shoulder (Hallman *et al.*, 2015) and in the low back region (Gupta *et al.*, 2015; Hallman *et al.*, 2016). However, results of the present investigation found no significant differences between the prevalence of musculoskeletal symptoms in any of the body regions when comparing the workers with lower sitting time and the workers experiencing higher sitting time, which is not in agreement with the previously mentioned studies. The same was reported regarding the physical activity level of the blue-collar workers of this sample, as no statistical differences in the prevalences of musculoskeletal symptoms were found between the workers within the three categories described in the IPAQ for any of the body regions. These results are also not in line with the study conducted by Haukka, Ojajärvi, Takala, Viikari-Juntura, and Leino-Arjas (2012) who concluded that high levels of physical activity have a significantly positive effect on musculoskeletal pain or discomfort in employees (Haukka *et al.*, 2012). Similarly, Van Amelsvoort *et al.* (2006) demonstrated a significantly less sickness absence by musculoskeletal disorders in active workers in their leisure time than sedentary ones, which suggests the benefits of physical activity. However, it should be noted that this investigation was conducted in blue-collar workers, which in itself represents a higher physical demanding job, with less sitting time (Schreuder *et al.*, 2008). Thus, the classification of an employee as physically active can be mostly influenced by their labor activity and not by the time they spend actually doing leisure time physical activity. This can explain the difference between the results presented here and the results of previous investigations, as they only quantified the leisure time physical activity of the assessed workers (Haukka *et al.*, 2012). This assumption is confirmed by previous studies conducted in blue-collar workers, where it was quantified the actual time they spent in occupational and leisure time physical activity, being demonstrated that this population spends higher periods of time walking and less time sitting during the work day than in the leisure time (Hallman *et al.*, 2015), which can have influenced the present results for these variables.

As previously mentioned, musculoskeletal pain in the working population can have multifactorial causes, in which age, high BMI, occupational stress and work-related physical factors such as heavy loading, twisting movements of the trunk, working with a flexed spinal posture or the hands above shoulders level, seem to play an important role (Aghilinejad *et al.*, 2012; Chen *et al.*, 2005; da Costa & Vieira, 2010; de Zwart *et al.*, 1997; Goswami *et al.*, 2016; Kerr *et al.*, 2001; Miranda *et al.*, 2001). Due to high morbidity rates, the importance of preventive measures has been increasingly emphasized in the last years. In that sense, several authors have performed different workplace physical activity interventions aiming to decrease workers BMI, sick leave, medical costs; to increase employees physical activity levels; and to prevent new episodes of pain and physical discomfort, and consequently, continuous or new musculoskeletal disorders among workers, demonstrating significantly positive effects (Christensen *et al.*, 2011; Kellett *et al.*, 1991; Kietrys *et al.*, 2007; Moreira-Silva, Santos, Abreu, & Mota, 2014; Tveito *et al.*, 2004; Tveito & Eriksen, 2009; Zebis *et al.*, 2011).

For future studies, when studying the associations between physical activity and musculoskeletal pain among the working-age population, researchers should also address the influence of others factors which are strongly related to pain, such as individual, psychosocial and work-related factors (Chen *et al.*, 2005; Kerr *et al.*, 2001; Miranda *et al.*, 2001; Osilla *et al.*, 2012; Widanarko *et al.*, 2011). More research with prospective design is needed in order to achieve more reliable information about the true effects of physical activity not only on the musculoskeletal health, but also for example on the delay of the influence of the individual factors, like aging; and on the psychosocial factors, as workplace physical activity interventions can promote and improve workers relationships/interaction.

A limitation of this study is that the sample, although not small, may have not been large enough to reveal significant associations between some explored variables. Moreover, the assessment of physical activity levels was done using an instrument that does not differentiate the physical activity performed in the leisure context from the physical activity in a labor context.

## 5. CONCLUSIONS

The population of the study (i.e., employees in job groups with high physical demands) is well documented to have a high risk of physical deterioration, as shown by the high 12-month and 7-day prevalence of musculoskeletal symptoms.

In the present sample, individual factors like age, gender and BMI were associated with a higher prevalence of musculoskeletal symptoms in different body regions, but lifestyle-related factors such as physical activity level and sitting time seem not to be associated with the prevalence of musculoskeletal symptoms in the blue-collar workers of this sample.

If proven effective, specific interventions in this job group can provide scientifically meaningful information for the development of public health promotion strategies for employees at higher risk for physical deterioration, such as blue-collar workers. This knowledge can be beneficial for occupational health professionals, companies and employees in these job groups. Despite the difficulties, an epidemiological perspective is needed since both musculoskeletal symptoms and physical exercise concern vast populations.

## ACKNOWLEDGEMENTS

This paper is an extended version of the book chapter "Prevalence of musculoskeletal symptoms in blue-collar workers: association with gender and physical activity level", published in the book: Occupational and Environmental Safety and Health (Publisher: Springer) (Moreira-Silva, Azevedo, Rodrigues, Seixas, & Mota, 2019).

## REFERENCES

- Aghilinejad, M., Choobineh, A., Sadeghi, Z., Nouri, M., & Ahmadi, A. B. (2012). Prevalence of musculoskeletal disorders among Iranian steel workers. *Iranian Red Crescent Medical Journal*, 14(4), 198-203.
- Andersen, J. H., Haahr, J. P., & Frost, P. (2007). Risk factors for more severe regional musculoskeletal symptoms: A two-year prospective study of a general working population. *Arthritis & Rheumatism*, 56(4), 1355-1364.
- Bureau of Labor Statistics Occupational Safety and Health. (2002). Lost-Worktime Injuries And Illnesses: Characteristics And Resulting Days Away From Work. Bureau of Labor Statistics. Retrieved from United States Department of Labor: <http://www.bls.gov/iif/>
- Bureau of Labor Statistics Occupational Safety and Health. (2012). Occupational Safety and Health Definitions. Bureau of Labor Statistics. Retrieved from United States Department of Labor: <http://www.bls.gov/iif/oshdef.htm>
- Chen, C., Perry, P., Yang, Y., & Yang, C. (2017). Decent work in the Chinese apparel industry: Comparative analysis of blue-collar and white-collar garment workers. *Sustainability*, 9(8), 1344-1361.
- Chen, W. Q., Yu, I. T. S., & Wong, T. W. (2005). Impact of occupational stress and other psychosocial factors on musculoskeletal pain among Chinese offshore oil installation workers. *Occupational And Environmental Medicine*, 62(4), 251-256.
- Christensen, J., Faber, A., Ekner, D., Overgaard, K., Holtermann, A., & Sogaard, K. (2011). Diet, physical exercise and cognitive behavioral training as a combined workplace based intervention to reduce body weight and increase physical capacity in health care workers - a randomized controlled trial. *BMC Public Health*, 11(1), 671-682. doi:10.1186/1471-2458-11-671



- Craig, C. L., Marshall, A. L., Sjoström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., & Oja, P. (2003). International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*, 35(8), 1381-1395.
- da Costa, B. R., & Vieira, E. R. (2010). Risk factors for work-related musculoskeletal disorders: a systematic review of recent longitudinal studies. *American Journal Of Industrial Medicine*, 53(3), 285-323.
- De Magalhães, J. P., Curado, J., & Church, G. M. (2009). Meta-analysis of age-related gene expression profiles identifies common signatures of aging. *Bioinformatics*, 25(7), 875-881.
- de Zwart, B. C. H., Broersen, J. P. J., Frings-Dresen, M. H. W., & Van Dijk, F. J. H. (1997). Musculoskeletal complaints in the Netherlands in relation to age, gender and physically demanding work. *International archives of occupational and environmental health*, 70(5), 352-360.
- Ghasemkhani, M., Mahmudi, E., & Jabbari, H. (2008). Musculoskeletal symptoms in workers. *International Journal of Occupational Safety and Ergonomics*, 14(4), 455-462.
- Goetzel, R. Z., & Ozminkowski, R. J. (2008). The health and cost benefits of work site health-promotion programs. *Annu Rev Public Health*, 29, 303-323.
- Goswami, S., Dasgupta, S., Samanta, A., Talukdar, G., Chanda, A., Ray Karmakar, P., . . . Chakrabarti, A. (2016). Load handling and repetitive movements are associated with chronic low back pain among Jute Mill Workers in India. *Pain research and treatment*, 2016(2016), 8.
- Gupta, N., Christiansen, C. S., Hallman, D. M., Korshøj, M., Carneiro, I. G., & Holtermann, A. (2015). Is objectively measured sitting time associated with low back pain? A cross-sectional investigation in the NOMAD study. *PLoS One*, 10(3), 1-18.
- Gupta, N., Hallman, D. M., Mathiassen, S. E., Aadahl, M., Jørgensen, M. B., & Holtermann, A. (2016). Are temporal patterns of sitting associated with obesity among blue-collar workers? A cross sectional study using accelerometers. *BMC Public Health*, 16:148. doi:DOI 10.1186/s12889-016-2803-9
- Hallman, D. M., Gupta, N., Heiden, M., Mathiassen, S. E., Korshøj, M., Jørgensen, M. B., & Holtermann, A. (2016). Is prolonged sitting at work associated with the time course of neck-shoulder pain? A prospective study in Danish blue-collar workers. *BMJ open*, 6(11), e012689.
- Hallman, D. M., Gupta, N., Mathiassen, S. E., & Holtermann, A. (2015). Association between objectively measured sitting time and neck-shoulder pain among blue-collar workers. *International archives of occupational and environmental health*, 88(8), 1031-1042.
- Hanebuth, D., Meinel, M., & Fischer, J. E. (2006). Health-related quality of life, psychosocial work conditions, and absenteeism in an industrial sample of blue-and white-collar employees: a comparison of potential predictors. *Journal of Occupational and Environmental Medicine*, 48(1), 28-37.
- Haukka, E., Ojajärvi, A., Takala, E.-P., Viikari-Juntura, E., & Leino-Arjas, P. (2012). Physical workload, leisure-time physical activity, obesity and smoking as predictors of multisite musculoskeletal pain. A 2-year prospective study of kitchen workers. *Occup Environ Med*, 69(7), 485-492.
- Holtermann, A., Jørgensen, M. B., Gram, B., Christensen, J. R., Faber, A., Overgaard, K., . . . Søgaard, K. (2010). Worksite interventions for preventing physical deterioration among employees in job-groups with high physical work demands: Background, design and conceptual model of FINALE. *BMC Public Health*, 10, 120-131.
- Hussain, T. (2004). Musculoskeletal symptoms among truck assembly workers. *Occupational Medicine*, 54(8), 506-512.
- Kellett, K. M., Kellett, D. A., & Nordholm, L. A. (1991). Effects of an exercise program on sick leave due to back pain. *Physical Therapy*, 71(4), 283-293.
- Kerr, M. S., Frank, J. W., Shannon, H. S., Norman, R. W., Wells, R. P., Neumann, W. P., . . . Group, O. U. B. P. S. (2001). Biomechanical and psychosocial risk factors for low back pain at work. *American Journal of Public Health*, 91(7), 1069.
- Kietrys, D. M., Galper, J. S., & Verno, V. (2007). Effects of At-work Exercises on Computer Operators. *Work: Journal of Prevention, Assessment & Rehabilitation*, 28(1), 67-75.
- Kuorinka, I. J., B. Kilbom, A. Vinterberg, H. Biering-Sørensen, F. Andersson, G. Jørgensen, K. (1987). Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Applied Ergonomics*, 18(3), 233-237. doi:10.1016/0003-6870(87)90010-x
- Manjunatha, R., Kiran, D., & Thankappan, K. R. (2011). Sickness absenteeism, morbidity and workplace injuries among iron and steel workers-A cross sectional study from Karnataka, Southern India. *The Australasian medical journal*, 4(3), 144-147.
- Mesquita, C., Ribeiro, J. C., & Moreira, P. (2010). Portuguese version of the standardized Nordic musculoskeletal questionnaire: cross cultural and reliability. *J Public Health* (2010), 18, 461-466. doi:10.1007/s10389-010-0331-0

- Miranda, Viikari-Juntura, E., Punnett, L., & Riihimäki, H. (2008). Occupational loading, health behavior and sleep disturbance as predictors of low-back pain. *Scandinavian Journal Of Work, Environment & Health*, 34(6), 411-419.
- Miranda, H., Viikari-Juntura, E., Martikainen, R., Takala, E. P., & Riihimäki, H. (2001). Physical exercise and musculoskeletal pain among forest industry workers. *Scandinavian Journal of Medicine & Science in Sports*, 11(4), 239-246. doi:10.1034/j.1600-0838.2001.110408.x
- Moreira-Silva, I., Azevedo, J., Rodrigues, S., Seixas, A., & Mota, J. (2019). Prevalence of Musculoskeletal Symptoms in Blue-Collar Workers: Association with Gender and Physical Activity Level. In *Occupational and Environmental Safety and Health* (pp. 467-475): Springer.
- Moreira-Silva, I., Santos, R., Abreu, S., & Mota, J. (2014). The effect of a physical activity program on decreasing physical disability indicated by musculoskeletal pain and related symptoms among workers: a pilot study. *International Journal of Occupational Safety and Ergonomics*, 20(1), 55-64.
- Ng, Y. G., Tamrin, S. B. M., Yik, W. M., Yusoff, I. S. M., & Mori, I. (2013). The prevalence of musculoskeletal disorder and association with productivity loss: a preliminary study among labour intensive manual harvesting activities in oil palm plantation. *Industrial health*, 52, 78-85.
- Osilla, K. C., Van, K. B., Schnyer, C., Larkin, J. W., Eibner, C., & Mattke, S. (2012). Systematic review of the impact of worksite wellness programs. *The American journal of managed care*, 18(2), e68-81.
- Schreuder, K. J., Roelen, C. A. M., Koopmans, P. C., & Groothoff, J. W. (2008). Job demands and health complaints in white and blue collar workers. *Work*, 31(4), 425-432.
- Serranheira, F. (2007). Lesões Musculo Esqueleticas ligadas com o Trabalho: Que Metodos de Avaliação do Risco? Retrieved from Universidade Nova de Lisboa:
- Sjögrene, T., Nissinen, K. J., Järvenpää, S. K., Ojanen, M. T., Vanharanta, H., & Mälkiä, E. A. (2006). Effects of a workplace physical exercise intervention on the intensity of low back symptoms in office workers: A cluster randomized controlled cross-over design. *Journal of Back & Musculoskeletal Rehabilitation*, 19(1), 13-24.
- Tinubu, B. M., Mbada, C. E., Oyeyemi, A. L., & Fabunmi, A. A. (2010). Work-Related Musculoskeletal Disorders among Nurses in Ibadan, South-west Nigeria: a cross-sectional survey. *BMC Musculoskeletal Disorders*, 11(1), 12-20. doi:10.1186/1471-2474-11-12
- Tveito, T., Hysing, M., & Eriksen, H. R. (2004). Low back pain interventions at the workplace: a systematic literature review. *Occupational Medicine*, 54(1), 3-13.
- Tveito, T. H., & Eriksen, H. R. (2009). Integrated health programme: A workplace randomized controlled trial. *Journal of Advanced Nursing*, 65(1), 110-119. doi:10.1111/j.1365-2648.2008.04846.x
- Van Amelsvoort, L. G. P. M., Spigt, M. G., Swaen, G. M. H., & Kant, I. (2006). Leisure time physical activity and sickness absenteeism; a prospective study. *Occupational Medicine*, 56(3), 210-212.
- Viestar, L., Verhagen, E. A., Hengel, K. M. O., Koppes, L. L., van der Beek, A. J., & Bongers, P. M. (2013). The relation between body mass index and musculoskeletal symptoms in the working population. *BMC Musculoskeletal Disorders*, 14(1), 238-246.
- Wang, Y., Tuomilehto, J., Jousilahti, P., Antikainen, R., Mähönen, M., Katzmarzyk, P. T., & Hu, G. (2010). Occupational, Commuting, and Leisure-Time Physical Activity in Relation to Heart Failure Among Finnish Men and Women. *Journal of the American College of Cardiology*, Vol. 56, No. 14. doi:10.1016
- WHO. (1995). Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. World Health Organization. Retrieved from Geneva:
- Widanarko, B., Legg, S., Stevenson, M., Devereux, J., Eng, A., Cheng, S., . . . Pearce, N. (2011). Prevalence of musculoskeletal symptoms in relation to gender, age, and occupational/industrial group. *International Journal of Industrial Ergonomics*, 41(5), 561-572.
- Widanarko, B., Legg, S., Stevenson, M., Devereux, J., & Jones, G. (2013). Prevalence of low back symptoms and its consequences in relation to occupational group. *American Journal Of Industrial Medicine*, 56(5), 576-589.
- World Health Organization, W. (2019). Body Mass Index - BMI. Retrieved from <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>
- Yildirim, Y., Gunay, S., & Karadibak, D. (2014). Identifying factors associated with low back pain among employees working at a package producing industry. *Journal of Back and Musculoskeletal Rehabilitation*, 27(1), 25-32.
- Zebis, M. K., Andersen, L. L., Pedersen, M. T., Mortensen, P., Andersen, C. H., Pedersen, M. M., . . . Mortensen, O. S. (2011). Implementation of neck/shoulder exercises for pain relief among industrial workers: a randomized controlled trial. *BMC Musculoskeletal Disorders*, 12(1), 205-213.